

Generalized Euler-Lagrange equations for variational problems with scale derivatives*

Ricardo Almeida
ricardo.almeida@ua.pt

Delfim F. M. Torres
delfim@ua.pt

Department of Mathematics
University of Aveiro
3810-193 Aveiro, Portugal

Abstract

We obtain several Euler-Lagrange equations for variational functionals defined on a set of Hölder curves. The cases when the Lagrangian contains multiple scale derivatives, depends on a parameter, or contains higher-order scale derivatives are considered.

Mathematics Subject Classification: 49K05, 26B05, 39A12.

Keywords: scale calculus, Euler-Lagrange equations, higher-order scale derivatives, non-differentiability.

1 Introduction

In 1992 L. Nottale introduced the theory of scale-relativity without the hypothesis of space-time differentiability [6, 7]. A rigorous mathematical foundation to Nottale's scale-relativity theory was then given by J. Cresson in 2005 [3, 4]. Roughly speaking, the calculus of variations developed in [3] cover sets of non differentiable curves, by substituting the classical derivative by a new complex operator, known as the *scale derivative*. Here we proceed with the theory started in [3] and continued in [1, 5], by presenting several Euler-Lagrange equations in the class of Hölderian curves and generalizing the previous results. For the necessary terminology and motivation to the study of such non-differentiable calculus of variations we refer the reader to [1, 2, 3, 6].

The paper is organized as follows. In Section 2 we review the necessary notions of scale calculus. Our results are then given in Section 3: (i) in §3.1 we prove the Euler-Lagrange equation for functionals defined by multiple scale derivatives; (ii) in §3.2 we deduce the Euler-Lagrange equation with a dependence on a complex parameter ξ ; (iii) finally, in §3.3 we characterize the extremals of variational functionals containing higher-order scale derivatives.

*Submitted on 03-Aug-2009; accepted for publication 16-March-2010; in *Letters in Mathematical Physics*.

2 Preliminaries

In what follows, function $f \in C^0$ and $\epsilon > 0$ is a real number. The ϵ -left and ϵ -right quantum derivatives are defined by the formulas

$$\Delta_\epsilon^- f(x) = -\frac{f(x-\epsilon) - f(x)}{\epsilon} \quad \text{and} \quad \Delta_\epsilon^+ f(x) = \frac{f(x+\epsilon) - f(x)}{\epsilon},$$

respectively. The ϵ scale derivative of f at x is given by

$$\square_\epsilon f(x) = \frac{1}{2}(\Delta_\epsilon^+ f(x) + \Delta_\epsilon^- f(x)) - i\frac{1}{2}(\Delta_\epsilon^+ f(x) - \Delta_\epsilon^- f(x)).$$

Theorem 1 ([3]). *Let $f, g \in C^0$ and $\epsilon > 0$. Then,*

$$\square_\epsilon(f \cdot g) = \square_\epsilon f \cdot g + f \cdot \square_\epsilon g + i\frac{\epsilon}{2}(\square_\epsilon f \square_\epsilon g - \square_\epsilon f \square_\epsilon g - \square_\epsilon f \square_\epsilon g - \square_\epsilon f \square_\epsilon g),$$

where $\square_\epsilon(*)$ is the complex conjugate of $\square_\epsilon(*)$.

The space of curves under consideration is the space of Hölderian curves with Hölder exponent α . More precisely, given a real number $\alpha \in (0, 1)$ and a sufficiently small real parameter ϵ , $0 < \epsilon \ll 1$, we define

$$C_\epsilon^\alpha(a, b) = \{y : [a - \epsilon, b + \epsilon] \rightarrow \mathbb{R} \mid y \in H^\alpha\}.$$

In [3] variational functionals of the type $\Phi(y) = \int_a^b f(x, y(x), \square_\epsilon y(x)) dx$ are studied in the class $y \in C_\epsilon^\alpha(a, b)$. As variation curves, it is considered those of the following type: let β be a positive real satisfying condition

$$\beta \geq (1 - \alpha)\mathbb{1}_{]0, 1/2[} + \alpha\mathbb{1}_{[1/2, 1]},$$

and $h \in C_\epsilon^\beta(a, b)$ be such that $h(a) = 0 = h(b)$. A curve of the form $y + h$ is called a variation of y . One says that Φ is differentiable on $C_\epsilon^\alpha(a, b)$ if, for all curves $y \in C_\epsilon^\alpha(a, b)$ and for all variations $y + h$,

$$\Phi(y + h) - \Phi(y) = F_y(h) + R_y(h),$$

where F_y is a linear operator on the space $C_\epsilon^\beta(a, b)$ and $R_y(h) = O(h^2)$. A curve y is an extremal for Φ on $C_\epsilon^\beta(a, b)$ if $[F_y(h)]_\epsilon = 0$ for all $\epsilon > 0$ and all $h \in C_\epsilon^\beta(a, b)$.

Theorem 2 ([3]). *The curve y is an extremal for Φ on $C_\epsilon^\beta(a, b)$ if and only if*

$$[\partial_2 f(x, y(x), \square_\epsilon y(x)) - \square_\epsilon(\partial_3 f)(x, y(x), \square_\epsilon y(x))]_\epsilon = 0$$

for every $\epsilon > 0$.

3 Main Results

We obtain several generalizations of Theorem 2: Theorem 4 of §3.1 coincides with Theorem 2 in the particular case $n = 1$; Theorem 6 of §3.2 coincides with Theorem 2 in the particular case when the Lagrangian L does not depend on the parameter ξ ; Theorem 8 of §3.3 coincides with Theorem 2 in the particular case when the Lagrangian L does not depend on $\square_\epsilon^2 y(x)$ and ξ .

3.1 Euler-Lagrange equation for multiple scale derivatives

Let $\epsilon := (\epsilon_1, \dots, \epsilon_n) \in \mathbb{R}^n$ be a vector such that $0 < \epsilon_k \ll 1$ for all $k \in \{1, \dots, n\}$. For continuous functions f let

$$\triangle_{\epsilon_k}^\sigma f(x) = \sigma \frac{f(x + \sigma \epsilon_k) - f(x)}{\epsilon_k}, \quad \sigma = \pm.$$

We define the ϵ_k scale derivative of f at x by the rule

$$\square_{\epsilon_k} f(x) = \frac{1}{2} (\triangle_{\epsilon_k}^+ f(x) + \triangle_{\epsilon_k}^- f(x)) - i \frac{1}{2} (\triangle_{\epsilon_k}^+ f(x) - \triangle_{\epsilon_k}^- f(x)).$$

Let $\alpha \in (0, 1)$ be a real, $\epsilon_M := \max\{\epsilon_1, \dots, \epsilon_n\}$, and

$$C_{\epsilon_M}^\alpha(a, b) = \{y : [a - \epsilon_M, b + \epsilon_M] \rightarrow \mathbb{R} \mid y \in H^\alpha\}.$$

We consider the following type of functionals: $\Phi : C_{\epsilon_M}^\alpha(a, b) \rightarrow \mathbb{C}$ of form

$$\Phi(y) = \int_a^b L(x, y(x), \square_{\epsilon_1} y(x), \dots, \square_{\epsilon_n} y(x)) dx, \quad (1)$$

where $L : \mathbb{R}^2 \times \mathbb{C}^n \rightarrow \mathbb{C}$ is a given C^1 function, called the Lagrangian. As in [3], we assume the Lagrangian L to satisfy

$$\|DL(x, y(x), \square_{\epsilon_1} y(x), \dots, \square_{\epsilon_n} y(x))\| \leq C$$

for all x and $\epsilon_1 > 0, \dots, \epsilon_n > 0$, where C is a positive constant, D denotes the differential, and $\|\cdot\|$ is a norm for matrices.

Definition 1. We say that Φ , as in (1), is differentiable on $C_{\epsilon_M}^\alpha(a, b)$ if for all curves $y \in C_{\epsilon_M}^\alpha(a, b)$ and for all variations $y + h$, $h \in C_{\epsilon_M}^\beta(a, b)$, the equality

$$\Phi(y + h) - \Phi(y) = F_y(h) + R_y(h)$$

holds with F_y a linear operator on the space $C_{\epsilon_M}^\beta(a, b)$ and $R_y(h) = O(h^2)$.

Let us determine the expression of $F_y(h)$. To simplify notation, let $u := (x, y, \square_{\epsilon_1} y, \dots, \square_{\epsilon_n} y)$. Then,

$$\begin{aligned} \Phi(y + h) - \Phi(y) &= \int_a^b [L(x, y + h, \square_{\epsilon_1} y + \square_{\epsilon_1} h, \dots, \square_{\epsilon_n} y + \square_{\epsilon_n} h) - L(x, y, \square_{\epsilon_1} y, \dots, \square_{\epsilon_n} y)] dx \\ &= \int_a^b [\partial_2 L(u) \cdot h + \partial_3 L(u) \cdot \square_{\epsilon_1} h + \dots + \partial_{n+2} L(u) \cdot \square_{\epsilon_n} h] dx + O(h^2). \end{aligned}$$

Integrating by parts (see Theorem 1), we obtain:

$$\begin{aligned} F_y(h) &= \int_a^b \left[\partial_2 L(u) \cdot h + \square_{\epsilon_1} (\partial_3 L(u) \cdot h) - \square_{\epsilon_1} (\partial_3 L(u)) \cdot h - i \frac{\epsilon_1}{2} \Sigma_{\epsilon_1} (\partial_3 L(u), h) \right. \\ &\quad \left. + \dots + \square_{\epsilon_n} (\partial_{n+2} L(u) \cdot h) - \square_{\epsilon_n} (\partial_{n+2} L(u)) \cdot h - i \frac{\epsilon_n}{2} \Sigma_{\epsilon_n} (\partial_{n+2} L(u), h) \right] dx, \end{aligned}$$

where

$$\Sigma_{\epsilon_k}(p, q) = \square_{\epsilon_k} p \square_{\epsilon_k} q - \square_{\epsilon_k} p \square_{\epsilon_k} q - \square_{\epsilon_k} p \square_{\epsilon_k} q - \square_{\epsilon_k} p \square_{\epsilon_k} q.$$

Therefore,

$$\begin{aligned} F_y(h) &= \int_a^b [\partial_2 L(u) - \square_{\epsilon_1}(\partial_3 L(u)) - \dots - \square_{\epsilon_n}(\partial_{n+2} L(u))] \cdot h \, dx \\ &\quad + \int_a^b [\square_{\epsilon_1}(\partial_3 L(u) \cdot h) + \dots + \square_{\epsilon_n}(\partial_{n+2} L(u) \cdot h)] \, dx \\ &\quad - i \int_a^b \left[\frac{\epsilon_1}{2} \Sigma_{\epsilon_1}(\partial_3 L(u), h) + \dots + \frac{\epsilon_n}{2} \Sigma_{\epsilon_n}(\partial_{n+2} L(u), h) \right] \, dx. \end{aligned}$$

We just proved the following result:

Theorem 3. *For all $\epsilon_1 > 0, \dots, \epsilon_n > 0$, the functional Φ defined by (1) is differentiable, and*

$$\begin{aligned} F_y(h) &= \int_a^b [\partial_2 L(u) - \square_{\epsilon_1}(\partial_3 L(u)) - \dots - \square_{\epsilon_n}(\partial_{n+2} L(u))] \cdot h \, dx \\ &\quad + \int_a^b [\square_{\epsilon_1}(\partial_3 L(u) \cdot h) + \dots + \square_{\epsilon_n}(\partial_{n+2} L(u) \cdot h)] \, dx \\ &\quad - i \int_a^b \left[\frac{\epsilon_1}{2} \Sigma_{\epsilon_1}(\partial_3 L(u), h) + \dots + \frac{\epsilon_n}{2} \Sigma_{\epsilon_n}(\partial_{n+2} L(u), h) \right] \, dx. \end{aligned}$$

Definition 2. *Let $a(\epsilon)$, $\epsilon = (\epsilon_1, \dots, \epsilon_n)$, be a real or complex valued function. We denote by $[\cdot]_\epsilon$ the linear operator such that*

$$a(\epsilon) - [a(\epsilon)]_\epsilon \xrightarrow{\epsilon \rightarrow 0} 0 \quad \text{and} \quad [a(\epsilon)]_\epsilon = 0 \quad \text{if} \quad \lim_{\epsilon \rightarrow 0} a(\epsilon) = 0.$$

Definition 3. *A curve y is an extremal for Φ on $C_{\epsilon_M}^\beta(a, b)$ if $[F_y(h)]_\epsilon = 0$ for all $\epsilon_1 > 0, \dots, \epsilon_n > 0$ and $h \in C_{\epsilon_M}^\beta(a, b)$.*

We now prove an Euler-Lagrange type equation for functionals of type (1).

Theorem 4. *A curve y is an extremal for Φ defined by (1) on $C_{\epsilon_M}^\beta(a, b)$ if and only if*

$$\left[\partial_2 L(u) - \sum_{k=1}^n \square_{\epsilon_k}(\partial_{k+2} L)(u) \right]_\epsilon = 0 \quad (2)$$

for all $\epsilon_1 > 0, \dots, \epsilon_n > 0$, where $u = (x, y(x), \square_{\epsilon_1} y(x), \dots, \square_{\epsilon_n} y(x))$.

Proof. Let $k \in \{1, \dots, n\}$. Then, for all $t \in [a, b]$

$$\sup_{x \in [t, t + \sigma \epsilon_k]} |\partial_{k+2} L(x, y(x), \square_{\epsilon_1} y(x), \dots, \square_{\epsilon_n} y(x))| \leq C_k \epsilon_m^{\alpha-1},$$

for some constant C_k , where $\epsilon_m := \min\{\epsilon_1, \dots, \epsilon_n\}$. Therefore, as ϵ goes to zero, the quantities

$$\int_a^b \square_{\epsilon_k}(\partial_{k+2}L(u) \cdot h) dx \quad \text{and} \quad \epsilon_k \int_a^b \Sigma_{\epsilon_k}(\partial_{k+2}L(u), h) dx$$

also go to zero (cf. [3, Lemma 3.2]). Applying the bracket operator $[\cdot]_\epsilon$ to $F_y(h)$, the formula (2) follows from the fundamental lemma of the calculus of variations. $\square$

In [1] we study the isoperimetric problem in the scale calculus context, considering integral constraints containing scale derivatives. There, with the help of an appropriate auxiliary function, we prove a necessary condition for extremals. We now extend the main result of [1] to functionals containing multiple scale derivatives.

Definition 4. Let $\Phi(y) = \int_a^b L(x, y(x), \square_{\epsilon_1}y(x), \dots, \square_{\epsilon_n}y(x)) dx$ and $\Psi(y) = \int_a^b g(x, y(x), \square_{\epsilon_1}y(x), \dots, \square_{\epsilon_n}y(x)) dx$ be two functionals on $C_{\epsilon_M}^\alpha(a, b)$. A curve y is called an extremal of Φ subject to the constraint $\Psi(y) = c$, $c \in \mathbb{C}$, if for all $m \in \mathbb{N}$ and all variations $\hat{y} = y + \sum_{k=1}^m h_k$, where $(h_k)_{1 \leq k \leq m} \in C_{\epsilon_M}^\beta(a, b)$ are such that $\Psi(\hat{y}) = c$, one has $[F_y(h_k)]_\epsilon = 0$ for all $\epsilon_1 > 0, \dots, \epsilon_n > 0$ and all $k \in \{1, \dots, m\}$.

Using the techniques in the proof of Theorem 4 and in [1, Theorem 4], the following result can be easily obtained.

Theorem 5. Suppose that $y \in C_{\epsilon_M}^\alpha(a, b)$ is an extremal for the functional Φ on $C_{\epsilon_M}^\beta(a, b)$ subject to the constraint $\Psi(y) = c$, $c \in \mathbb{C}$. If

1. y is not an extremal for Ψ ;
2. both limits

$$\lim_{\epsilon \rightarrow 0} \max_{x \in [a, b]} |\partial_2 L - \square_{\epsilon_1}(\partial_3 L) - \dots - \square_{\epsilon_n}(\partial_{n+2} L)|$$

and

$$\lim_{\epsilon \rightarrow 0} \max_{x \in [a, b]} |\partial_2 g - \square_{\epsilon_1}(\partial_3 g) - \dots - \square_{\epsilon_n}(\partial_{n+2} g)|$$

are finite along y ;

then there exists $\lambda \in \mathbb{C}$ such that

$$[\partial_2 K - \square_{\epsilon_1}(\partial_3 K) - \dots - \square_{\epsilon_n}(\partial_{n+2} K)]_\epsilon = 0$$

holds along the curve y , where $K = L - \lambda g$.

3.2 Dependence on a parameter

We now characterize the extremals in the case when the variational functional depends on a complex parameter ξ . We consider functionals of the form

$$\Phi(y, \xi) = \int_a^b L(x, y(x), \square_\epsilon y(x), \xi) dx, \quad (3)$$

where $(y, \xi) \in C_\epsilon^\alpha(a, b) \times \mathbb{C}$. We say that Φ is differentiable if for all $(y, \xi) \in C_\epsilon^\alpha(a, b) \times \mathbb{C}$ and for all $(h, \delta) \in C_\epsilon^\beta(a, b) \times \mathbb{C}$ one has

$$\Phi(y + h, \xi + \delta) - \Phi(y, \xi) = F_{(y, \xi)}(h, \delta) + R_{(y, \xi)}(h, \delta),$$

where $F_{(y, \xi)}$ is a linear operator on the space $C_\epsilon^\beta(a, b) \times \mathbb{C}$ and $R_{(y, \xi)}(h, \delta) = O(|(h, \delta)|^2)$. With similar calculations as done before, we deduce:

$$\begin{aligned} & \int_a^b \left[L(x, y + h, \square_\epsilon y + \square_\epsilon h, \xi + \delta) - L(x, y, \square_\epsilon y, \xi) \right]_\epsilon dx \\ &= \int_a^b [\partial_2 L(u) \cdot h + \partial_3 L(u) \cdot \square_\epsilon h + \partial_4 L(u) \cdot \delta]_\epsilon dx \\ &= \int_a^b [\partial_2 L(u) - \square_\epsilon(\partial_3 L(u))]_\epsilon \cdot h dx + \int_a^b [\partial_4 L(u)]_\epsilon \cdot \delta dx, \end{aligned}$$

where $u := (x, y(x), \square_\epsilon y(x), \xi)$. For $\delta = 0$ we obtain the Euler-Lagrange equation $[\partial_2 L(u) - \square_\epsilon(\partial_3 L(u))]_\epsilon = 0$ for all $\epsilon > 0$. For $h = 0$ we get $\int_a^b [\partial_4 L(u)]_\epsilon dx = 0$. In summary, we have:

Theorem 6. *The pair (y, ξ) is an extremal for Φ given by (3), i.e., $[F_{(y, \xi)}(h, \delta)]_\epsilon = 0$, if and only if*

$$\begin{cases} [\partial_2 L(u) - \square_\epsilon(\partial_3 L(u))]_\epsilon = 0 \\ \int_a^b [\partial_4 L(u)]_\epsilon dx = 0 \end{cases}$$

for all $\epsilon > 0$, where $u = (x, y(x), \square_\epsilon y(x), \xi)$.

Example 1. *Let Φ be given by the expression*

$$\Phi(y, \xi) = \int_{-1}^1 (\square_\epsilon y - \square_\epsilon |x|)^2 + (\xi x)^2 dx.$$

Then, $(y, \xi) = (|x|, 0)$ is an extremal of Φ :

$$[\partial_2 L(u) - \square_\epsilon(\partial_3 L(u))]_\epsilon = [-\square_\epsilon(2(\square_\epsilon y - \square_\epsilon |x|))]_\epsilon = 0$$

and

$$\int_{-1}^1 [\partial_4 L(u)]_\epsilon dx = \int_{-1}^1 [2\xi x^2]_\epsilon dx = 0.$$

Example 2. *Consider now the functional $\Phi(y, \xi) = \int_{-1}^1 (\xi \cdot \square_\epsilon y - \square_\epsilon |x|)^2 dx$, $\xi \in \mathbb{C}$. Similarly as in Example 1, it can be proved that $(y, \xi) = (|x|, 1)$ is an extremal of Φ . Observe that if we substitute y by $|x|$ and $\xi \in \mathbb{R}$ in Φ , simple calculations show that $\Phi(|x|, \xi) = 2(\xi - 1)^2(1 - \epsilon)$. This function has a global minimizer for $\xi = 1$.*

3.3 Higher-order Euler-Lagrange equation

Let $\alpha \in (0, 1)$, $\epsilon > 0$, $n \in \mathbb{N}$, and $f \in C^{n-1}[a, b]$ be a real valued function.

Definition 5. For $k = 1, \dots, n$ let

$$\Delta_\epsilon^{k,\sigma} f(x) = \sigma \frac{f^{(k-1)}(x + \sigma\epsilon) - f^{(k-1)}(x)}{\epsilon}, \quad \sigma = \pm.$$

We define the k th ϵ -scale derivative of f at x by

$$\square_\epsilon^k f(x) = \frac{1}{2} (\Delta_\epsilon^{k,+} f(x) + \Delta_\epsilon^{k,-} f(x)) - i \frac{1}{2} (\Delta_\epsilon^{k,+} f(x) - \Delta_\epsilon^{k,-} f(x)).$$

Consider variational functionals of the form

$$\Phi(y) = \int_a^b L(x, y(x), \square_\epsilon^1 y(x), \dots, \square_\epsilon^n y(x)) dx,$$

for curves y of class C^{n-1} and $y, \square_\epsilon^1 y, \dots, \square_\epsilon^{n-1} y \in C_\epsilon^\alpha(a, b)$. Observe that, as $\epsilon \rightarrow 0$, we obtain the standard functional of the calculus of variations with higher-order derivatives:

$$\Phi(y) = \int_a^b L(x, y(x), y'(x), \dots, y^{(n)}(x)) dx.$$

We study the case $n = 2$:

$$\Phi(y) = \int_a^b L(x, y(x), \square_\epsilon^1 y(x), \square_\epsilon^2 y(x)) dx. \quad (4)$$

Results for the general case are easily proved by induction. Let h be a function of class C^1 such that $h, \square_\epsilon^1 h \in C_\epsilon^\beta(a, b)$, $h(a) = 0 = h(b)$ and $h'(a) = 0 = h'(b)$. Observe that $\square_\epsilon^2 y = \square_\epsilon^1 y'$ and $\square_\epsilon^2 h = \square_\epsilon^1 h'$. Thus, by Theorem 1 and the standard integration by parts formula (here we are assuming that L and y are at least of class C^2),

$$\begin{aligned} & [\Phi(y + h) - \Phi(y)]_\epsilon \\ &= \int_a^b [L(x, y + h, \square_\epsilon^1 y + \square_\epsilon^1 h, \square_\epsilon^2 y + \square_\epsilon^2 h) - L(x, y, \square_\epsilon^1 y, \square_\epsilon^2 y)]_\epsilon dx \\ &= \int_a^b [\partial_2 L \cdot h + \partial_3 L \cdot \square_\epsilon^1 h + \partial_4 L \cdot \square_\epsilon^2 h]_\epsilon dx + O(h^2) \\ &= \int_a^b [\partial_2 L \cdot h + \partial_3 L \cdot \square_\epsilon^1 h + \partial_4 L \cdot \square_\epsilon^1 h']_\epsilon dx + O(h^2) \\ &= \int_a^b [\partial_2 L \cdot h - \square_\epsilon^1(\partial_3 L) \cdot h - \square_\epsilon^1(\partial_4 L) \cdot h']_\epsilon dx + O(h^2) \\ &= \int_a^b [\partial_2 L - \square_\epsilon^1(\partial_3 L) + (\square_\epsilon^1(\partial_4 L))']_\epsilon \cdot h dx + O(h^2). \end{aligned}$$

We just deduced the Euler-Lagrange equation for (4):

Theorem 7. *Let L be a Lagrangian of class C^2 , and Φ as in (4) be defined on the class C^2 of curves such that $y, \square_\epsilon^1 y \in C_\epsilon^\alpha(a, b)$. Function y is an extremal for Φ if and only if*

$$[\partial_2 L - \square_\epsilon^1(\partial_3 L) + (\square_\epsilon^1(\partial_4 L))']_\epsilon = 0$$

for all $\epsilon > 0$.

Remark 1. *In contrast with the classical theory of the calculus of variations for functionals containing second-order derivatives, where typically admissible functions are of class $C^4[a, b]$, here it is enough to work with C^2 curves.*

We can easily include the case when the functional depends on a complex parameter ξ , as was done in Section 3.2:

Theorem 8. *Let Φ be the functional defined by*

$$\Phi(y, \xi) = \int_a^b L(x, y(x), \square_\epsilon^1 y(x), \square_\epsilon^2 y(x), \xi) dx.$$

The pair (y, ξ) is an extremal for Φ if and only if

$$\begin{cases} [\partial_2 L - \square_\epsilon^1(\partial_3 L) + (\square_\epsilon^1(\partial_4 L))']_\epsilon = 0 \\ \int_a^b [\partial_5 L(u)]_\epsilon dx = 0 \end{cases}$$

for all $\epsilon > 0$, where $u = (x, y(x), \square_\epsilon^1 y(x), \square_\epsilon^2 y(x), \xi)$.

Acknowledgments

Work supported by the *Centre for Research on Optimization and Control* (CEOC) from the “Fundação para a Ciência e a Tecnologia” (FCT), cofinanced by the European Community Fund FEDER/POCI 2010.

References

- [1] R. Almeida and D. F. M. Torres, Hölderian variational problems subject to integral constraints, *J. Math. Anal. Appl.* **359** (2009), no. 2, 674–681. [arXiv:0807.3076](#)
- [2] J. Cresson, Scale calculus and the Schrödinger equation, *J. Math. Phys.* **44** (2003), no. 11, 4907–4938.
- [3] J. Cresson, Non-differentiable variational principles, *J. Math. Anal. Appl.* **307** (2005), no. 1, 48–64.
- [4] J. Cresson, Non-differentiable deformations of $\mathbb{R}^n$, *Int. J. Geom. Methods Mod. Phys.* **3** (2006), no. 7, 1395–1415.
- [5] J. Cresson, G. S. F. Frederico and D. F. M. Torres, Constants of motion for non-differentiable quantum variational problems, *Topol. Methods Nonlinear Anal.* **33** (2009), no. 2, 217–232. [arXiv:0805.0720](#)
- [6] L. Nottale, The theory of scale relativity, *Internat. J. Modern Phys. A* **7** (1992), no. 20, 4899–4936.
- [7] L. Nottale, The scale-relativity program, *Chaos Solitons Fractals* **10** (1999), no. 2-3, 459–468.